



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

investigation he has attempted to show quantitatively the influence of nutrition. In the two ferns investigated, he found that the development of the two sex organs could be controlled by certain concentrations of Knop's solution. In general, the number of antheridia decreases as the concentration decreases. In both species archegonia are formed only above 0.175 per cent Knop's solution, which was the optimum concentration for *Asplenium*; but 0.35 per cent was the optimum for *Osmunda*. In many of the prothallia of *Asplenium* the two sex organs appeared only successively. The prothallia of *Osmunda* remained almost completely sterile when grown in solutions lacking calcium or magnesium salts. Numerous combinations of conditions were used, with varying results, and the evidence all indicated that the production of sex organs is a response to factors in the environment.—J. M. C.

Indiana Academy.—The Proceedings of the Indiana Academy of Sciences for 1914 contains the following contributions of interest to botanists: An apparatus for aerating culture solutions, by PAUL WEATHERWAX; Antagonism of *B. fluorescens* and *B. typhosus* in culture, by P. A. TETRAULT; Notes upon the distribution of forest trees in Indiana, by STANLEY COULTER; Mosses of Monroe County, Indiana, by MILDRED NOTHNAGEL and F. L. PICKETT; A new enemy of the black locust, by GLENN CULBERTSON; A new leaf spot of *Viola cucullata*, by H. W. ANDERSON; Oat smut in Indiana, by F. J. PIPEL; Plants new or rare to Indiana, by C. C. DEAN; Some peculiarities in *Spirogyra dubia*, by PAUL WEATHERWAX; Report on corn pollination, by M. L. FISHER; Stomata of *Trillium nivale*, by F. M. ANDREWS; The primrose-leaved violet in White County, by L. M. HEIMLICH; Continuous rust propagation without sexual reproduction, by C. A. LUDWIG; Correlation of certain long-cycled and short-cycled rusts, by H. C. TRAVELBEE; Some species of *Nummularia* common in Indiana, by C. E. O'NEAL; The genus *Rosellinia* in Indiana, by G. B. RAMSEY; Some large botanical problems, by J. C. ARTHUR.—J. M. C.

A new genus of Bennettitales.—THOMAS³¹ has described a new genus of Bennettitales (*Williamsoniella*) based upon material obtained from mesozoic beds in Yorkshire. It is represented by buds, mature strobili ("flowers"), microsporophylls, and the ovulate portions of the strobili. The strobilus is bisporangiate and very small, with no ensheathing sterile bracts; 12–16 wedge-shaped microsporophylls, each bearing 4–6 synangia; sessile ovules, "very similar in external appearance to the interseminal scales"; and the sterile tip of the strobilus axis terminating in "a characteristic corona-like structure" (which suggested the specific name, *W. coronata*). In all probability these strobili were borne in the forks of dichotomously branching stems, whose leaves had been known as *Taeniopteris vittata*. A "flower bud," thought to belong to the same genus, is named *W. roseberriensis*. The marked features

³¹ THOMAS, H. H., On *Williamsoniella*, a new type of Bennettitalean flower. Phil. Trans. Roy. Soc. London B 207: 113–148. pls. 12–14. 1915.